

YÜK TERİMLERİ TABLOSU

Yük	\mathcal{L}, \mathcal{R}	Yük	\mathcal{L}, \mathcal{R}
	$\mathcal{L} = \mathcal{R} = \frac{ql^2}{4}$		$\mathcal{L} = \mathcal{R} = \frac{q}{4} \left[l^2 - a^2 \left(2 - \frac{a}{l} \right) \right]$
	$\mathcal{L} = \mathcal{R} = \frac{qa^2}{4} \left(2 - \frac{a}{l} \right)^2$ $\mathcal{R} = \frac{qa^2}{4} \left(2 - \frac{a^2}{l^2} \right)$		$\mathcal{L} = \mathcal{R} = \frac{l^2}{60} (8q_1 + 7q_2)$ $\mathcal{R} = \frac{l^2}{60} (7q_1 + 8q_2)$
	$\mathcal{L} = \mathcal{R} = \frac{qc}{8l} (3l^2 - c^2)$		$\mathcal{L} = \mathcal{R} = \frac{ql^2}{5}$
	$\mathcal{L} = \frac{qbc}{l^2} \left(l^2 - b^2 - \frac{c^2}{4} \right)$ $\mathcal{R} = \frac{qac}{l^2} \left(l^2 - a^2 - \frac{c^2}{4} \right)$		$\mathcal{L} = \frac{Pab(b+l)}{l^2}$ $\mathcal{R} = \frac{Pab(a+l)}{l^2}$
	$\mathcal{L} = \mathcal{R} = \frac{qa^2}{2l} (3l - 2a)$		$\mathcal{L} = \mathcal{R} = \frac{3}{8} Pl$
	$\mathcal{L} = \mathcal{R} = \frac{5}{32} ql^2$		$\mathcal{L} = \mathcal{R} = 3Pa \left(1 - \frac{a}{l} \right)$
	$\mathcal{L} = \mathcal{R} = \frac{ql^2}{60} \left(1 + \frac{b}{l} \right) \left(7 - 3 \frac{b^2}{l^2} \right)$ $\mathcal{R} = \frac{ql^2}{60} \left(1 + \frac{a}{l} \right) \left(7 - 3 \frac{a^2}{l^2} \right)$		$\mathcal{L} = \mathcal{R} = Pl \frac{n}{4} \left(1 - \frac{1}{n^2} \right)$ n: tek sayı
	$\mathcal{L} = \mathcal{R} = \frac{8}{60} ql^2$ $\mathcal{R} = \frac{7}{60} ql^2$		$\mathcal{L} = \mathcal{R} = Pl \frac{n}{4} \left(1 + \frac{1}{2n^2} \right)$ n: tek sayı
	$\mathcal{L} = \mathcal{R} = \frac{qa^2}{4l} (2l - a)$		$\mathcal{L} = M \left(1 - \frac{3b^2}{l^2} \right)$ $\mathcal{R} = M \left(\frac{3a^2}{l^2} - 1 \right)$